

Recommendations for EMS Clinicians and Medical First Responders

EMS clinician practices should be based on the most up-to-date COVID-19 clinical recommendations and information from appropriate public health authorities and EMS medical direction.

State and local EMS authorities may direct EMS clinicians to modify their practices as described below.

Patient assessment

- If PSAP call takers advise that the patient is suspected of having COVID-19, EMS clinicians should put on appropriate [PPE](#) before entering the scene. EMS clinicians should consider the signs, symptoms, and risk factors of COVID-19 (<https://www.cdc.gov/coronavirus/2019-nCoV/clinical-criteria.html>).
- If information about potential for COVID-19 has not been provided by the PSAP, EMS clinicians should exercise appropriate precautions when responding to any patient with signs or symptoms of a respiratory infection. Initial assessment should begin from a distance of at least 6 feet from the patient, if possible. Patient contact should be minimized to the extent possible until a facemask is on the patient. If COVID-19 is suspected, all [PPE](#) as described below should be used. If COVID-19 is not suspected, EMS clinicians should follow standard procedures and use appropriate PPE for evaluating a patient with a potential respiratory infection.
- A facemask should be worn by the patient for source control. If a nasal cannula is in place, a facemask should be worn over the nasal cannula. Alternatively, an oxygen mask can be used if clinically indicated. If the patient requires intubation, see below for additional precautions for aerosol-generating procedures.
- During transport, limit the number of providers in the patient compartment to essential personnel to minimize possible exposures.

Recommended Personal Protective Equipment (PPE)

- EMS clinicians who will directly care for a patient with possible COVID-19 infection or who will be in the compartment with the patient should follow Standard, Contact, and Airborne Precautions, including the use of eye protection. Recommended PPE includes:
 - A single pair of disposable patient examination gloves. Change gloves if they become torn or heavily contaminated,
 - Disposable isolation gown,
 - Respiratory protection (i.e., N-95 or higher-level respirator), and
 - Eye protection (i.e., goggles or disposable face shield that fully covers the front and sides of the face).
- Drivers, if they provide direct patient care (e.g., moving patients onto stretchers), should wear all recommended PPE. After completing patient care and before entering an isolated driver's compartment, the driver should remove and dispose of PPE and perform hand hygiene to avoid soiling the compartment.
 - If the transport vehicle does **not** have an isolated driver's compartment, the driver should remove the face shield or goggles, gown and gloves and perform hand hygiene. A respirator should continue to be used during transport.
- All personnel should avoid touching their face while working.
- On arrival, after the patient is released to the facility, EMS clinicians should remove and discard PPE and perform hand hygiene. Used PPE should be discarded in accordance with routine procedures.
- Other required aspects of Standard Precautions (e.g., injection safety, hand hygiene) are not emphasized in this document but can be found in the guideline titled [Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings](#).

Precautions for Aerosol-Generating Procedures

- If possible, consult with medical control before performing aerosol-generating procedures for specific guidance.
- In addition to the PPE described above, EMS clinicians should exercise caution if an aerosol-generating procedure (e.g., bag valve mask (BVM) ventilation, oropharyngeal suctioning, endotracheal intubation, nebulizer treatment, continuous positive airway pressure (CPAP), bi-phasic positive

airway pressure (biPAP), or resuscitation involving emergency intubation or cardiopulmonary resuscitation (CPR) is necessary.

- BVMs, and other ventilatory equipment, should be equipped with HEPA filtration to filter expired air.
- EMS organizations should consult their ventilator equipment manufacturer to confirm appropriate filtration capability and the effect of filtration on positive-pressure ventilation.
- If possible, the rear doors of the transport vehicle should be opened and the HVAC system should be activated during aerosol-generating procedures. This should be done away from pedestrian traffic.

EMS Transport of a PUI or Patient with Confirmed COVID-19 to a Healthcare Facility (including interfacility transport)

If a patient with an exposure history and signs and symptoms suggestive of COVID-19 requires transport to a healthcare facility for further evaluation and management (subject to EMS medical direction), the following actions should occur during transport:

- EMS clinicians should notify the receiving healthcare facility that the patient has an exposure history and signs and symptoms suggestive of COVID-19 so that appropriate infection control precautions may be taken prior to patient arrival.
- Keep the patient separated from other people as much as possible.
- Family members and other contacts of patients with possible COVID-19 should **not** ride in the transport vehicle, if possible. If riding in the transport vehicle, they should wear a facemask.
- Isolate the ambulance driver from the patient compartment and keep pass-through doors and windows tightly shut.
- When possible, use vehicles that have isolated driver and patient compartments that can provide separate ventilation to each area.
 - Close the door/window between these compartments before bringing the patient on board.
 - During transport, vehicle ventilation in both compartments should be on non-recirculated mode to maximize air changes that reduce potentially infectious particles in the vehicle.

- If the vehicle has a rear exhaust fan, use it to draw air away from the cab, toward the patient-care area, and out the back end of the vehicle.
- Some vehicles are equipped with a supplemental recirculating ventilation unit that passes air through HEPA filters before returning it to the vehicle. Such a unit can be used to increase the number of air changes per hour (ACH)
(<https://www.cdc.gov/niosh/hhe/reports/pdfs/1995-0031-2601.pdf>pdf icon).
- If a vehicle without an isolated driver compartment and ventilation must be used, open the outside air vents in the driver area and turn on the rear exhaust ventilation fans to the highest setting. This will create a negative pressure gradient in the patient area.
- Follow routine procedures for a transfer of the patient to the receiving healthcare facility (e.g., wheel the patient directly into an Airborne Infection Isolation Room).

Documentation of patient care

- Documentation of patient care should be done after EMS clinicians have completed transport, removed their PPE, and performed hand hygiene.
 - Any written documentation should match the verbal communication given to the emergency department providers at the time patient care was transferred.
- EMS documentation should include a listing of EMS clinicians and public safety providers involved in the response and level of contact with the patient (for example, no contact with patient, provided direct patient care). This documentation may need to be shared with local public health authorities.

Cleaning EMS Transport Vehicles after Transporting a PUI or Patient with Confirmed COVID-19

The following are general guidelines for cleaning or maintaining EMS transport vehicles and equipment after transporting a PUI:

- After transporting the patient, leave the rear doors of the transport vehicle open to allow for sufficient air changes to remove potentially infectious particles.
 - The time to complete transfer of the patient to the receiving facility and complete all documentation should provide sufficient air changes.
- When cleaning the vehicle, EMS clinicians should wear a disposable gown and gloves. A face shield or facemask and goggles should also be worn if splashes or sprays during cleaning are anticipated.
- Ensure that environmental cleaning and disinfection procedures are followed consistently and correctly, to include the provision of adequate ventilation when chemicals are in use. Doors should remain open when cleaning the vehicle.
- Routine cleaning and disinfection procedures (e.g., using cleaners and water to pre-clean surfaces prior to applying an EPA-registered, hospital-grade disinfectant to frequently touched surfaces or objects for appropriate contact times as indicated on the product's label) are appropriate for SARS-CoV-2 (the virus that causes COVID-19) in healthcare settings, including those patient-care areas in which aerosol-generating procedures are performed.
- Products with EPA-approved emerging viral pathogens claims are recommended for use against SARS-CoV-2. These products can be identified by the following claim:
 - “[Product name] has demonstrated effectiveness against viruses similar to SARS-CoV-2 on hard non-porous surfaces. Therefore, this product can be used against SARS-CoV-2 when used in accordance with the directions for use against [name of supporting virus] on hard, non-porous surfaces.”
 - This claim or a similar claim, will be made only through the following communications outlets: technical literature distributed exclusively to health care facilities, physicians, nurses and public health officials, “1-800” consumer information services, social media sites and company websites (non-label related). Specific claims for “SARS-CoV-2” will not appear on the product or master label.

- See [additional information about EPA-approved emerging viral pathogens claimsexternal icon](#).
- If there are no available EPA-registered products that have an approved emerging viral pathogen claim, products with label claims against human coronaviruses should be used according to label instructions.
- Clean and disinfect the vehicle in accordance with standard operating procedures. All surfaces that may have come in contact with the patient or materials contaminated during patient care (e.g., stretcher, rails, control panels, floors, walls, work surfaces) should be thoroughly cleaned and disinfected using an EPA-registered hospital grade disinfectant in accordance with the product label.
- Clean and disinfect reusable patient-care equipment before use on another patient, according to manufacturer's instructions.
- Follow standard operating procedures for the containment and disposal of used PPE and regulated medical waste.
- Follow standard operating procedures for containing and laundering used linen. Avoid shaking the linen.

Criteria to Guide Evaluation of PUI for COVID-19

Local health departments, in consultation with clinicians, should determine whether a patient is a PUI for COVID-2019. The CDC clinical criteria for COVID-19 PUIs have been developed based on available information about this novel virus, as well as what is known about [Severe Acute Respiratory Syndrome \(SARS\)](#) and [Middle East Respiratory Syndrome \(MERS\)](#). These criteria are subject to change as additional information becomes available.

Clinical features and epidemiologic risk		
Clinical Features	&	Epidemiologic Risk
Fever ¹ or signs/symptoms of lower respiratory illness (e.g. cough or shortness of breath)	AND	Any person, including healthcare workers ² , who has had close contact ³ with a laboratory-confirmed ⁴ COVID-19 patient within 14 days of symptom onset
Fever ¹ and signs/symptoms of a lower respiratory illness (e.g., cough or shortness of breath) requiring hospitalization	AND	A history of travel from affected geographic areas ⁵ (see below) within 14 days of symptom onset
Fever ¹ with severe acute lower respiratory illness (e.g., pneumonia, ARDS) requiring hospitalization and without alternative explanatory diagnosis (e.g., influenza) ⁶	AND	No source of exposure has been identified